

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

| CANDIDATE NAME | | |
|--------------------|---------------------------|-----------------------|
| CENTRE NUMBER | | CANDIDATE NUMBER |
| CHEMISTRY | | 0620/33 |
| Paper 3 (Extended | () | October/November 2011 |
| | | 1 hour 15 minutes |
| Candidates answe | er on the Question Paper. | |
| No Additional Mate | erials are required. | |

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in. Write in dark blue or black pen. You may use a pencil for any diagrams, graphs or rough working. Do not use staples, paper clips, highlighters, glue or correction fluid. DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions. A copy of the Periodic Table is printed on page 12.

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question.

| For Examiner's Use | | | |
|--------------------|--|--|--|
| 1 | | | |
| 2 | | | |
| 3 | | | |
| 4 | | | |
| 5 | | | |
| 6 | | | |
| 7 | | | |
| Total | | | |

This document consists of **11** printed pages and **1** blank page.



UNIVERSITY of CAMBRIDGE International Examinations www.sparkl.me

| (a) | (a) Choose an element from the Periodic Table to match each description. You may give either the name or the symbol. | | | | | | |
|-----|---|--|--|--|--|--|--|
| | (i) | It is the most reactive metal[1] | | | | | |
| | (ii) | It is the only non-metal which is a liquid at r.t.p [1] | | | | | |
| (| iii) | An isotope of this element is used as a fuel in nuclear reactors | | | | | |
| (| (iv) This Group VII element is a solid at r.t.p. [1 | | | | | | |
| | (v) This element is in Group V and Period 4 | | | | | | |
| (| (vi) This unreactive gas is used to fill lamps [| | | | | | |
| (b) | (b) Predict the formula of each of the following compounds. | | | | | | |
| | (i) germanium oxide | | | | | | |
| | (ii) tellurium bromide | | | | | | |
| (c) | c) Give the formula of each of the following ions. | | | | | | |
| | (i) | strontium | | | | | |
| | (ii) | fluoride[2] | | | | | |
| | | [Total: 10] | | | | | |

Use your copy of the Periodic Table to answer these questions.

1

www.sparkl.me

For Examiner's Use

| | 3 | |
|---------------------------|--|----------------------|
| | a complex carbohydrate, is a natural macromolecule or polymer. e formed from its monomer by condensation polymerisation. | For Examin Use |
| (a) (i) | Explain the terms: | |
| | monomer | |
| | | |
| | condensation polymerisation | |
| | [2] | |
| (ii) | Draw the structural formula of starch to include three monomer units. | |
| | Glucose, the monomer, can be represented as HOOH. | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | [3] | |
| wai | [3] arch can be hydrolysed to simple sugars by heating with dilute sulfuric acid or by rming with a dilute solution of saliva. The reaction can be catalysed by H ⁺ ions from acid or by the enzymes in saliva. | |
| wai | arch can be hydrolysed to simple sugars by heating with dilute sulfuric acid or by rming with a dilute solution of saliva. The reaction can be catalysed by H ⁺ ions from | |
| wai the | arch can be hydrolysed to simple sugars by heating with dilute sulfuric acid or by rming with a dilute solution of saliva. The reaction can be catalysed by H ⁺ ions from acid or by the enzymes in saliva. | |
| wai the (i) | arch can be hydrolysed to simple sugars by heating with dilute sulfuric acid or by rming with a dilute solution of saliva. The reaction can be catalysed by H ⁺ ions from acid or by the enzymes in saliva. What is an enzyme? | |
| wai the | arch can be hydrolysed to simple sugars by heating with dilute sulfuric acid or by rming with a dilute solution of saliva. The reaction can be catalysed by H ⁺ ions from acid or by the enzymes in saliva. What is an enzyme? | |
| wai the (i) | arch can be hydrolysed to simple sugars by heating with dilute sulfuric acid or by rming with a dilute solution of saliva. The reaction can be catalysed by H ⁺ ions from acid or by the enzymes in saliva. What is an enzyme? [1] Explain why, if the saliva/starch mixture is heated above 70 °C, the hydrolysis stops. [1] The complete acid-catalysed hydrolysis of starch forms only glucose. The partial acid-catalysed hydrolysis of starch forms a mixture of sugars which | |
| wai the (i) (ii) | arch can be hydrolysed to simple sugars by heating with dilute sulfuric acid or by rming with a dilute solution of saliva. The reaction can be catalysed by H ⁺ ions from acid or by the enzymes in saliva. What is an enzyme? (1] Explain why, if the saliva/starch mixture is heated above 70 °C, the hydrolysis stops. (1] The complete acid-catalysed hydrolysis of starch forms only glucose. | |
| wai the (i) (ii) | arch can be hydrolysed to simple sugars by heating with dilute sulfuric acid or by rming with a dilute solution of saliva. The reaction can be catalysed by H ⁺ ions from acid or by the enzymes in saliva. What is an enzyme? [1] Explain why, if the saliva/starch mixture is heated above 70 °C, the hydrolysis stops. [1] The complete acid-catalysed hydrolysis of starch forms only glucose. The partial acid-catalysed hydrolysis of starch forms a mixture of sugars which | |
| wai the (i) (ii) | arch can be hydrolysed to simple sugars by heating with dilute sulfuric acid or by training with a dilute solution of saliva. The reaction can be catalysed by H* ions from acid or by the enzymes in saliva. What is an enzyme? [1] Explain why, if the saliva/starch mixture is heated above 70 °C, the hydrolysis stops. [1] The complete acid-catalysed hydrolysis of starch forms only glucose. The partial acid-catalysed hydrolysis of starch forms a mixture of sugars which includes glucose. Describe how you could identify the different sugars in this mixture. | |
| wai the (i) (ii) | arch can be hydrolysed to simple sugars by heating with dilute sulfuric acid or by rming with a dilute solution of saliva. The reaction can be catalysed by H ⁺ ions from acid or by the enzymes in saliva. What is an enzyme? [1] Explain why, if the saliva/starch mixture is heated above 70 °C, the hydrolysis stops. [1] The complete acid-catalysed hydrolysis of starch forms only glucose. The partial acid-catalysed hydrolysis of starch forms a mixture of sugars which | |

0620/33/O/N/11

| For | |
|------------|--|
| Examiner's | |
| | |



| The rea | ions in this question are all examples of photochemical reactions. | | | | |
|----------------|--|--|--|--|--|
| (a) Exp | plain the phrase photochemical reaction. | | | | |
| | | | | | |
| | | | | | |
| cor | iny millions of years ago, the Earth's atmosphere was rich in carbon dioxide and intained negligible amounts of oxygen. After the appearance of green plant-like cteria, the proportions of these two gases in the atmosphere changed. | | | | |
| (i) | What are the approximate percentages of these two gases in the atmosphere now? | | | | |
| | carbon dioxide =[1] | | | | |
| | oxygen =[1] | | | | |
| (ii) | Explain how the green plant-like bacteria changed the composition of the atmosphere. | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| (c) The | e reduction of silver(I) bromide to silver is the basis of film photography. | | | | |
| | $2AaBr \rightarrow 2Aa + Br$ | | | | |

6

For

Examiner's Use

 $2AgBr \rightarrow 2Ag + Br_2$ white black

An opaque object is placed on a piece of paper coated with silver(I) bromide which is then exposed to a bright light. The light is switched off and the opaque object removed.



5

- 6 Nickel is a transition element.
 - (a) Predict three differences in the chemical properties of nickel and barium.

.....[3]

(b) Nickel ores are converted into nickel(II) oxide. This can be reduced to impure nickel by heating with carbon. The nickel is purified by the following reversible reaction.

 $Ni(s) + 4CO(g) \rightleftharpoons Ni(CO)_4(g)$

nickel carbonyl

(i) Impure nickel is heated at 60 °C. The forward reaction occurs.

 $Ni(s) + 4CO(g) \rightarrow Ni(CO)_4(g)$ impure

The nickel carbonyl, a gas, moves into a hotter chamber at 200 °C. The backward reaction occurs and the nickel carbonyl decomposes.

 $Ni(CO)_4(g) \rightarrow Ni(s) + 4CO(g)$ pure

Is the forward reaction exothermic or endothermic? Give a reason for your answer.

(ii) Explain why the forward reaction is favoured by an increase in pressure. [2]

(iii) Suggest what happens to the impurities.

......[1]

For

Examiner's Use (iv) Suggest another method of refining nickel. Give a brief description of the method which you have suggested. A labelled diagram is acceptable.

For Examiner's Use

[4]

[Total: 12]

www.sparkl.me

- The alkenes are a series of unsaturated hydrocarbons. They have the general molecular 7 Examiner's formula $C_n H_{2n}$.
 - (a) Deduce the molecular formula of an alkene which has a relative molecular mass of 126. Show your working.

.....

(b) The structural formula of propene is drawn below.



(i) Draw a diagram showing the arrangement of the valency electrons in one molecule of this covalent compound. Use x to represent an electron from an atom of carbon. Use o to represent an electron from an atom of hydrogen.

(ii) Draw the structure of the polymer formed from propene

[2]

[3]

For

Use

Use the data in the table to show that the following reaction is exothermic. н Н Н Н Н Н —Н =Ċ \rightarrow H-Н C -H н Н Н Н Н [3] (c) This question is concerned with some of the addition reactions of but-1-ene. (i) Name the product formed when but-1-ene reacts with water.[1] (ii) Complete the equation. $CH_3 - CH_2 - CH = CH_2 + Br_2 \rightarrow \dots$ [2] (iii) Deduce the formula of the compound which reacts with but-1-ene to form 1-iodobutane. [Total: 14]

(iii) Bond energy is the amount of energy, in kJ, which must be supplied to break one

bond energy

in kJ/mol +436

+610

+346

+415

For Examiner's Use

0620/33/O/N/11

www.sparkl.me

10

bond

 $\frac{H-H}{C=C}$

C-C

C-H

mole of the bond.

BLANK PAGE

11

www.sparkl.me

| | 0 | 4 Helium | 20 Neon 10 Ad Argon 18 Argon | 84 Krypton 36 | 131 Xe S4 Xenon | Radon 86 | 175 Lutetium 71 Lawrendum 103 |
|---|-------|------------|--|--|--|-----------------------------------|---|
| | M | | 19 9 Fluorine 35.5 C1 C1 C1 | 80 Br Bromine | 127 I fodine 53 | At Astatine 85 | 173 Yb 70 70 Nobelium 102 |
| | > | | 16 B Oxygen 32 32 Sulfur 16 | 79 Se Selenium 34 | 128 Te llurium 52 | Polonium 84 | 169 Tmulium 69 Mendelevium 101 |
| | | | 14 Nitrogen 31 Phosphorus 15 | 75 AS Arsenic 33 | 122 Sb 51 209 | Bismuth 83 | 167 Ertium 68 Fm Fm 100 |
| | ≥ | - | 6 Carbon 6 28 28 14 Silicon | 73 Ge Germanium 32 | 119 Sn 50 207 | 82 Lead | 165 Hohmuum 67 Einsteinium 99 |
| | ≡ | | 11 B Boron 27 Aluminium 13 | 70 Ga Gallium 31 | 115 Ind Indium 204 | TT Thallium 81 | 162 Dysprosium 66 Cf Californium |
| ents | | | | 65 Zn 30 ^{Zinc} | 112 Cadmium 48 201 | Mercury 08 | 159 Tb 5 Berkelium 97 |
| The Periodic Table of the Elements Group | | | | 64 Cu ^{Copper} | 108 AG 47 197 | Au Gold | 157 Gd Gd 64 64 Curium 96 |
| ble of th oup | Group | | | 59 Nickel 28 | 106 Palladium 195 | Platinum 78 | 152 Eu 63 Americium 95 |
| iodic Ta Gr | | | | 59 CO 27 | 103 Rh odium 45 192 | Ir Indium 77 | 150 Samartum 62 Plutonium 94 |
| The Per | | Hydrogen 1 | | 56 Fe Iron 26 | 101 Ru thenium 44 190 | Osmium 76 | 61 Reprunium 93 89 93 |
| | | | | 55 Mn ^{Manganese} 25 | Technetium 43 186 | Rhenium 75 | 144 Neodymium 60 238 238 Uanium |
| | | | | 52 Cr Chromium 24 | 96 Mo Molybdenum 42 184 | Tungsten 74 | 141 Praseodymium 59 Protactinium 91 |
| | | | | 51 V Vanadium 23 | 93 Niobium 41 | Tantalum 73 | 140 Centum 58 232 232 232 1hortum 90 |
| | | | | 48 Titanium 22 | 91 Zr Zirconium 40 178 | Hathium 72 | nic mass bol |
| | | | | 45 Sc Scandium 21 | 89 Vttrium 39 139 | Lanthanum 57 * | 89 1 0id series 1 I series a = relative atomic mass X = atomic symbol b = proton (atomic) number |
| | = | | 9 Be Beryllium 4 24 Ng Magnesium 12 | 40 Calcium 20 | 88 Strontium 38 137 | Barium 56 Barium 226 Radium | <pre>*58-71 Lanthanoid series *58-71 Lanthanoid series 190-103 Actinoid series Key key b b b b c c a a a relative a x = atomics b b b c c a b b c b c a b b c b c a b c b c</pre> |
| | - | | 7 Lithium 23 23 23 23 23 23 23 23 23 23 23 23 23 | 39 K Potassium 19 | 85 Rb ^{Rubidium} 37 | | *58-71 L ⁸⁷ *58-71 L ² 190-103 / Key |

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared wh reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwitting publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand na Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

12